

THE JOURNAL OF IMMUNOLOGY

Manuscripts should be typewritten and carefully revised before submission and should be sent to Dr. Arthur F. Coca, Pearl River, New York.

THE JOURNAL OF IMMUNOLOGY is issued monthly. Two volumes a year are issued.

Twenty-five reprints, without covers, of articles will be furnished to contributors free of cost when ordered in advance. A table showing cost of reprints, with an order slip, is sent with proof.

Published monthly, beginning January, 1929. (From 1916-1925, and in 1928 Volumes I-X, inclusive, and Vol. XV published bi-monthly. Two volumes per year were published during 1926-1927.)

The subscription price is \$5.00 per volume. For countries outside the Postal Union, add 50 cents a volume. Two volumes ordered at the same time, \$9.00; overseas, \$10.00. The subscriptions of members of the American Association of Immunologists are included in the membership dues and are handled by the Association's Treasurer, Dr. Arthur F. Coca, Pearl River, New York. All other subscriptions should be sent to the publishers.

Claims for copies lost in the mails must be received within 30 days, domestic (90 days, foreign) of the date of issue. Changes of address must be received at least two weeks in advance of issue.

Correspondence concerning business matters should be addressed to The Williams & Wilkins Company, Publishers of Scientific Journals and Books, Mount Royal and Guilford Avenues, Baltimore, U. S. A.

New subscriptions and renewals are entered to begin with the first issue of the current volume. Should any issue of the current volume be out of print at the time the subscription order is received, the pro-rata value of such numbers will be credited to the next volume, and the renewal invoice therefore adjusted accordingly.

Subscriptions should be renewed promptly—To avoid a break in your series, subscriptions should be renewed promptly. The publishers cannot guarantee to supply back issues on belated renewals.

Subscriptions, new or renewal, should be sent to the Publishers or to Agents listed below.

AGENTS

For Argentina and Uruguay: Beutelspacher y Cia, Sarmiento 815, Buenos Aires.

For Australia: Angus & Robertson, Limited, 89-95 Castlereagh Street, Sydney.

For Belgium: Henri Lamertin, 58 Rue Coudenberg, Bruxelles.

For the British Empire, except Australia and Canada: Baillière, Tindall & Cox, 8 Henrietta St., Covent Garden, W C 2, London, England.

For Canada: Wm. Dawson & Sons, Ltd., 91 Queen Street East, Toronto.

For China: Commercial Press Ltd., Paoshan Road, Shanghai, China.

For Denmark: H. Hagerup's Boghandel, Gothersgade 30, Kjobenhavn.

For France: Emile Bougault, 48 Rue des Ecoles, Paris.

For Germany: B. Westermann Co., Inc., Zimmerstrasse 35-41, Berlin SW-68.

For Holland: Scheltema & Holkema, Rokin 74-76, Amsterdam.

For Japan and Korea: Maruzen Company, Ltd. (Maruzen-Kabushiki-Kaisha), 6 Nihonbashi Tori-Nichome, Tokyo; Fukuoka, Osaka, Kyoto, and Sendai, Japan.

For Spain: Ruiz Hermanos, Plaza de Santa Ana, 13, Madrid.

THE WILLIAMS & WILKINS COMPANY

Publishers of Scientific Books and Periodicals

BALTIMORE, U. S. A.

PUBLISHERS OF: *Medicine, Journal of Urology, Journal of Pharmacology and Experimental Therapeutics, American Journal of Tropical Medicine, Journal of Immunology, Journal of Industrial Hygiene, Quarterly Review of Biology, Journal of Bacteriology, Annals of the Pickett-Thomson Research Laboratory, Chemical Reviews, Soil Science, Social Forces, Journal of Comparative Psychology, Mental Measurement Monographs, Occupational Therapy and Rehabilitation, Child Development, The American Journal of Clinical Pathology, Journal of Physical Chemistry, Journal of the Biological Photographic Association, Philosophy of Science, Medico-Legal and Criminological Review.*



Peptones for Bacteriological Use

This group of peptones is used expressly for preparing bacteriological culture media. The nitrogen content of the peptones is readily available for bacterial metabolism, permitting rapid and luxuriant growth of the inoculated organisms.

Bacto-Peptide

Bacto-Peptide is the standard peptide for all routine bacteriological uses. It is recommended as an ingredient of culture media in "Standard Methods." It is completely soluble in the usual concentrations and has a reaction of pH 7.0 in solution.

Proteose-Peptide

Proteose-Peptide is an excellent nutriment for the more fastidious pathogenic bacteria. Culture media prepared with this peptide are well suited for these organisms. The longevity of the organisms is prolonged and their virulence is maintained for considerable periods of time in media containing Proteose-Peptide.

Neopeptide

Neopeptide is free from the toxic elements which prevent or inhibit the development of the more delicate bacteria. Such organisms as the pneumococcus, which is notoriously difficult to propagate, will grow luxuriantly in media prepared with this peptide.

Bacto-Tryptone

Bacto-Tryptone is an excellent peptide for use in detecting the production of indol by bacteria. When Bacto-Tryptone is used, reliable and accurate results are obtained after relatively short periods of incubation. Bacto-Tryptone is also a very useful nutriment for many of the more fastidious bacteria.

Specify "DIFCO"

THE TRADE NAME OF THE PIONEERS

In the Research and Development of Bacto-Peptide and Dehydrated Culture Media

DIFCO LABORATORIES

INCORPORATED

DETROIT, MICHIGAN